arXiv:1601.04454v1 [math.AC] 18 Jan 2016

# LEFSCHETZ PROPERTIES FOR ARTINIAN GORENSTEIN ALGEBRAS PRESENTED BY QUADRICS

RODRIGO GONDIM* AND GIUSEPPE ZAPPALÀ

Abstract. We introduce a family of standard bigraded binomial Artinian Gorenstein algebras, whose combinatoric structure characterizes the ones presented by quadrics. These algebras provide, for all socle degree grater than two and in sufficiently large codimension with respect to the socle degree, counter-examples to Migliore-Nagel conjectures, see [MN1] and [MN2]. One of them predicted that Artinian Gorenstein algebras presented by quadratics should satisfy the weak Lefschetz property. We also prove a generalization of a Hessian criterion for the Lefschetz properties given by Watanabe, see [Wa1] and [MW], which is our main tool to control the Weak Lefschetz property.

## 1. Introduction

A standard graded $\mathbb{K}$ algebra is said to be presented by quadrics if it is isomorphic to the quotient of a polynomial ring over $\mathbb{K}$ by a homogeneous ideal generated by quadratic forms. Also called quadratic algebras, they are related with Koszul algebras and Grobner basis, see for example [Co]. From a more geometric point of view quadratic ideals appear as homogeneous ideals of very positive embeddings of any smooth projective varieties. As pointed out in [MN2], Artinian Gorenstein algebras presented by quadrics are also related to Eisenbud-Green-Harris conjectures motivated by the Cayley-Bacharach theorem. The Lefschetz properties have attracted a great deal of attention over the years, in a number of different settings including Commutative Algebra itself, Algebraic Geometry and Combinatorics, see [St, St2, MN1, HMMNWW, HMNW]. The present work lies in the border of these three areas.

More recently, in [MN2] the authors studied Artinian Gorenstein algebras presented by quadrics, they presented some constructions of such algebras and described their possible Hilbert vectors. In [MN1] and [MN2] the authors proposed two conjectures, which, from now on, we will call Migliore-Nagel conjectures. These conjectures inspired the present paper.

**Conjecture 1.1. (Migliore-Nagel injective Conjecture)** *For any Artinian Gorenstein algebra presented by quadrics, defined over a field $\mathbb{K}$ of characteristic zero, and of socle degree at least three, there exists $L \in A_1$, such that, the multiplication map $\bullet L : A_1 \to A_2$ is injective.*

**Conjecture 1.2. (Migliore-Nagel WLP Conjecture)** *Any Artinian Gorenstein algebra presented by quadrics, over a field $\mathbb{K}$ of characteristic zero, has the Weak Lefschetz Property. That is, there exists $L \in A_1$ such that all the maps $\bullet L : A_i \to A_{i+1}$ have maximal rank.*

In [MN2] the authors proved the WLP-conjecture for complete intersection of quadratic forms and presented computational evidence for the conjectures in low codimension. We

*Partially supported by the CAPES postdoctoral fellowship, Proc. BEX 2036/14-2.

want to stress the fact that as soon as the codimension increases with respect to the socle degree surprising phenomena begin to appear.

We recall that the Lefschetz properties for standard graded Artinian $\mathbb{K}$ algebra are algebraic abstractions introduced by Stanley in [St], motivated by the Hard Lefschetz Theorem on the cohomology rings of smooth irreducible complex projective varieties endowed with the euclidean topology, see for example the survey [La] for the theorem and [Ru] for an overview. The Poincaré duality for these cohomology rings inspired the definition of Poincaré duality algebras which, in this context, is equivalent to the Gorenstein hypothesis, for more details see, for example [MW] and [Ru]. In [Wa2] and [MW] the authors used Macaulay-Matlis duality in characteristic zero to present the Artinian Gorenstein algebra as $A = Q/\operatorname{Ann}_Q(f)$ where $f \in R = \mathbb{K}[x_1, \ldots, x_n]$ a polynomial ring and $Q = \mathbb{K}[X_1, \ldots, X_N]$ the associated ring of differential operators. They introduced a Higher Hessian determinantal criterion using a basis of $A_k$ to identify Strong Lefschetz elements $L \in A_1$. In [Go] this criterion was used by the first author to contruct explict examples of Artinian Gorenstein algebras that do not satisfy the Strong Lefschetz property and/or the Weak Lefschetz property, producing forms with vanishing Hessian of higher order.

Hypersurfaces with vanishing (first) Hessian is a classical subject that traces back to Hesse's wrong claim, see [He], that a form has vanihing Hessian if and only if its partial derivatives are linearly independent. Gordan-Noether in the fundamental paper [GN] proved a criterion to the vanishing of the Hessian, see Proposition 2.11. They showed that Hesse's claim is true for at most four variables and produced series of counterexamples for at least five variables, these counterexamples exaust all the examples in five variables. A modern proof of this fact can be found in [GR] while a very detailed account on the subject appears in [Ru]. Perazzo in [Pe] studied cubic hypersurfaces with vanishing Hessian in at most seven variables and found a canonial form that motivated our definition of Perazzo hypersurfaces also in degree $d \geq 3$, see Definition 2.12. More recently [CRS], [Lo], [GR], [GRu], [Wa2] and [Ru] treated different aspects of the subject, sometimes called Gordan-Noether Theory.

In [GRu], which contain results from the first author's thesis, Perazzo's work has been revisited from a modern point of view, re-obtaining the classification of cubics with vanishing Hessian in $\mathbb{P}^N$ for $N \leq 6$ and describing new examples in higher dimension. An important example of cubic with vanishing Hessian in $\mathbb{P}^7$, is given by a general tangent section of the secant variety of the Segre variety $\mathbb{P}^2 \times \mathbb{P}^2 \to \operatorname{Seg}(2,2) \subset \mathbb{P}^8$, see [GRu, p. 803, Example 6]. From the algebraic point of view, see Example 4.8, it provides a standard graded Artinian Gorenstein algebra presented by quadrics which does not have the WLP, being a counter example for both Migliore-Nagel conjectures. This example was the starting point of our work.

Our strategy to construct standard graded Artinian Gorenstein algebras presented by quadrics is to deal with the simplest ones, which are those whose ideal contains the complete intersection $(x_1^2, \ldots, x_n^2)$. This assumption forces all monomials that occur in $f$ to be square free. As a matter of fact we deal with bihomogeneous forms of bidegree $(1, d-1)$ of special type, here called of monomial square free type, see Definition 2.7, and note the similarity with Perazzo polynomial. To any bihomogeneous form of monomial square free type we can associate in a bijective way a pure simplicial complex whose combinatoric structure determine a set of generators of the annihilator ideal, see Theorem 3.1. This combinatoric object also characterizes when the associated algebra is presented by quadrics, see Theorem 3.3, wich

summarizes the main results of the work. Inspired by the famous Turan Theorem, see [Tu], that characterizes maximal graphs not containing a complete subgraph $K_l$, we introduce a simplicial complex, here called Turan complex, whose associated algebra is always presented by quadrics and such that, in sufficient large codimension and arbitrary degree the first multiplication map $\bullet L : A_1 \to A_2$ is not injective for all $L \in A_1$. Furthermore, for a particular subfamily, which occurs in very large codimension with respect to the socle degree, the Hilbert vector of $A$ is not even submodal in the first step, that is $\dim A_1 > \dim A_2$.

We now describe the contents of the paper in more detail. In the first section we recall the basic definitions and constructions of standard graded Artinian Gorenstein algebras, we deal also with the bigradedd case which is of particular interest, we recall the Lefschetz properties, Macaulay-Matlis duality and the hessian criterion given in [Wa1] and [MW]. The main result of the first section is a generalization of the Hessian criterion to mixed Hessians, see Theorem 2.10. We also recall classical results about forms with vanishing Hessian that will be useful in the sequel.

The second section is devoted to the main results and constructions. Theorem 3.1 describes the annihilator of a standard bigraded form of bidegree $(1, d-1)$ of monomial square free type, showing that it is a binomial ideal whose generators are closely connected with the combinatoric of the associated simplicial complex. Theorem 3.3 is the main result. It characterizes when such algebras are presented by quadrics. In this section we introduce the Turan complex and as Corollary 3.9, we produce counterexamples to both Migliore-Nagel conjectures in any socle degree and sufficient large codimension. It is surprising the fact that if the codimension is very large with respect to the socle degree, the Hilbert vector of the Turan algebras, that are quadratic and binomial, is non unimodal in the first step, an unexpected and very surprising fact.

In the last section we deal with an inductive construction, see Corollary 4.4. In the second subsection we give a complete description of bigraded algebras of monomial square free type of socle bidegree $(1, 2)$ that are presented by quadrics (see Theorem 4.11). In Corollary 4.12 we show that for all codimension $r \geq 8$ there are algebras of socle degree 3 presented by quadrics that does not satisfy the WLP. Furthermore the bound is sharp by Corollary4.7. For higher odd degree we prove the existence of algebras presented by quadrics that do not satisfy the WLP in all codimensions $r \geq d + 5$ (see Corollary 4.14). The bound is relatively sharp by the result for $d = 3$. In the last subsection we study the case of even socle degree, showing the strong difference with the odd socle degree case. We prove the existence of algebras of socle degree 4 and of codimension $r \geq 16$ that are presented by quadrics but not having the WLP, see Corollary 4.17. In the similar way we deal with the even socle degree in general (see Corollary 4.19). In even socle degree we are not able to give sharp bounds even in socle degree 4. The problem is that, since there is not a classification of quartics with vanishing Hessian in $\mathbb{P}^N$ for $N \geq 5$, there is not an analog of Corollary 4.7.

## 2. Lefschetz properties, Gordan-Noether theory and the Hessian criteria

2.1. **The Lefschetz Properties and Macaulay-Matlis duality.** Let $\mathbb{K}$ be an infinite field and $R = \mathbb{K}[x_1, \ldots, x_n]$ be the polynomial ring in $n$ indeterminates.

**Definition 2.1.** Let $A$ be a standard graded $\mathbb{K}$-algebra. We say that $A$ is presented by quadrics if $A \simeq R/I$, where $R = \mathbb{K}[x_1, \ldots, x_n]$ and the homogeneous ideal $I$ has a set of generators formed by quadratic forms.

Let $A = R/I$ be an Artinian standard graded $R$-algebra. If $F \in R_d$ is a form, it induces a $\mathbb{K}$-vector spaces map $\varphi_{i,F} : A_i \to A_{i+d}$, defined by $\varphi_{i,F}(\alpha) = F\alpha$, for every $\alpha \in A_i$.

**Definition 2.2.** We say that $A$ has the *Weak Lefschetz property* (in short WLP) if there exists a linear form $L \in R_1$ such that $\operatorname{rk} \varphi_{i,L} = \min\{\dim_{\mathbb{K}} A_i, \dim_{\mathbb{K}} A_{i+1}\}$, for every $i$.

**Definition 2.3.** We say that $A$ has the *Strong Lefschetz property* (in short SLP) if there exists a linear form $L \in R_1$ such that $\operatorname{rk} \varphi_{i,L^d} = \min\{\dim_{\mathbb{K}} A_i, \dim_{\mathbb{K}} A_{i+d}\}$, for every $i$ and $d$.

**Definition 2.4.** Let $A = R/I$ be an Artinian standard graded $R$-algebra. If $R = \mathbb{K}[x_1, \ldots, x_n]$ and $I_1 = 0$, then $n$ is said to be the codimension of $A$. If $A_d \neq 0$ and $A_i = 0$ for all $i > d$, then $d$ is called the socle degree of $A$. The Hilbert vector of $A$ is $h_A = \operatorname{Hilb}(A) = (1, h_1, h_2, \ldots, h_d)$, where $h_k = \dim A_k$. We say that $h_A$ is unimodal if there exists $k$ such that $1 \leq h_1 \leq \ldots \leq h_k \geq h_{k+1} \geq h_d$.

**Remark 2.5.** We recall that an Artinian algebra $A = \bigoplus_{i=0}^{d} A_i$, $A_d \neq 0$, is a *Gorenstein algebra* if $\dim_{\mathbb{K}} A_d = 1$ and the bilinear pairing

$$A_i \times A_{d-i} \to A_d$$

inducted by the multiplication is perfect for $0 \leq i \leq d$. So we have an isomorphism $A_i \simeq \operatorname{Hom}_{\mathbb{K}}(A_{d-i}, A_d)$ for $i = 0, \ldots, d$. In particular $\dim_{\mathbb{K}} A_i = \dim_{\mathbb{K}} A_{d-i}$, for $i = 0, \ldots, d$. Moreover for every $L \in R_1$, $\operatorname{rank} \varphi_{i,L} = \operatorname{rank} \varphi_{d-i-1,L}$, for $0 \leq i \leq d$.

Since $A$ as a $R$-module is generated in degree 0, if $\varphi_{i,L}$ is surjective, then $\varphi_{j,L}$ is surjective for every $j \geq i$. Therefore, if $A$ is a Gorenstein Artinian algebra, if $\varphi_{i,L}$ is injective, then $\varphi_{j,L}$ is injective for every $j \leq i$. Consequently a Gorenstein Artinian algebra $A = \bigoplus_{i=0}^{d} A_i$, $A_d \neq 0$, has the WLP iff there exists a linear form $L \in R_1$ such that $\varphi_{\frac{d}{2}-1,L}$ is injective when $d$ is even or $\varphi_{\frac{d-1}{2},L}$ is injective (or surjective) when $d$ is odd. We recall that for Gorenstein Artinian algebras the SLP is equivalent to the SLP in the narrow sense, that is $\varphi_{i,L^{d-2i}}$ to be an isomorphism. Notice also that the WLP implies the unimodality of the Hilbert vector of $A$. Unimodality in the Gorenstein case implies that $\dim A_{k-1} < \dim A_k$ for all $k \leq \frac{d}{2}$.

Now we assume that $\operatorname{char} \mathbb{K} = 0$. Let us regard the polynomial algebra $R$ as a module over the algebra $Q = \mathbb{K}[X_1, ..., X_n]$ via the identification $X_i = \partial/\partial x_i$. If $f \in R$ we set

$$\operatorname{Ann}_Q(f) = \{p(X_1, \ldots, X_n) \in Q \mid p(\partial/\partial x_1, \ldots, \partial/\partial x_n) f = 0\}.$$

By Macaulay-Matlis duality we have a bijection:

$$\begin{array}{ccc} \{\text{Homogeneous ideals of } R\} & \leftrightarrow & \{\text{Graded } Q \text{ modules}\} \\ \text{Ann}_Q(M) & \leftarrow & M \\ I & \rightarrow & I^{-1} \end{array}$$

Let $I \subset Q$ be an homogeneous ideal. It is well known that $A = Q/I$ is a Gorenstein standard graded Artinian algebra if and only if there exists a form $f \in R$ such that $I = \text{Ann}_Q(f)$ (for more details see, for example, [MW]).

In the sequel we always assume that $\text{char}(\mathbb{K}) = 0$, $A = Q/I$, $I = \text{Ann}_Q(f)$ and $I_1 = 0$. Sometimes we also assume that $\mathbb{K}$ is algebraically closed. All arguments work over $\mathbb{C}$.

We deal with standard bigraded Artinian Gorenstein algebras $A = \bigoplus_{i=0}^{d} A_i$, $A_d \neq 0$, with $A_k = \bigoplus_{i=0}^{k} A_{(i,k-1)}$, $A_{(d_1,d_2)} \neq 0$ for some $e_1, e_2$ such that $e_1 + e_2 = d$, we call $(d_1, d_2)$ the socle bidegree of $A$. Since $A_k^* \simeq A_{d-k}$ and since duality is compatible with direct sum, we get $A^*_{(i,j)} \simeq A_{(d_1-i,d_2-j)}$.

Let $R = \mathbb{K}[x_1, \ldots, x_n, u_1, \ldots, u_m]$ be the polynomial ring viewed as standard bigraded ring in the sets of variables $\{x_1, \ldots, x_n\}$ and $\{u_1, \ldots, u_m\}$ and let $Q = \mathbb{K}[X_1, \ldots, X_n, U_1, \ldots, U_m]$ be the associated ring of differential operators.

We want to stress that the bijection given by Macaulay-Matlis duality preserves bigrading, that is, there is a bijection:

$$\begin{array}{ccc} \{\text{Bihomogeneous ideals of } R\} & \leftrightarrow & \{\text{Bigraded } Q \text{ modules}\} \\ \text{Ann}_Q(M) & \leftarrow & M \\ I & \rightarrow & I^{-1} \end{array}$$

If $f \in R_{(d_1,d_2)}$ is a bihomogeneous polynomial of total degree $d = d_1 + d_2$, then $I = \text{Ann}_Q(f) \subset Q$ is a bihomogeneous ideal and $A = Q/I$ is a standard bigraded Artinian Gorenstein algebra of socle bidegree $(d_1, d_2)$ and codimension $r = m + n$ if we assume, without lost of generality, that $I_1 = 0$.

**Remark 2.6.** If $f \in R_{(d_1,d_2)}$ is a bihomogeneous polynomial of bidegree $(d_1, d_2)$, consider the associated bigraded algebra $A$ of socle bidegree $(d_1, d_2)$. Notice that for all $\alpha \in Q_{(i,j)}$ with $i > d_1$ or $j > d_2$ we get $\alpha(f) = 0$, therefore, in these conditions $I_{(i,j)} = Q_{(i,j)}$. As consequence, we have the following decomposition for all $A_k$:

$$A_k = \bigoplus_{i+j=k, i \leq d_1, j \leq d_2} A_{(i,j)}.$$

Furthermore, for $i < d_1$ and $j < d_1$, the evaluation map $Q_{i,j} \rightarrow A_{(d_1-i,d_2-j)}$ given by $\alpha \mapsto \alpha(f)$ provides the following short exact sequence:

$$0 \rightarrow I_{(i,j)} \rightarrow Q_{(i,j)} \rightarrow A_{(d_1-i,d_2-j)} \rightarrow 0.$$

One of our goals is to produce bigraded algebras of socle bidegree $(1, d-1)$ presented by quadrics. In order to achieve this objective we study the ideal of a particular family.

**Definition 2.7.** With the previous notation, all bihomogeneous polynomials of bidegree $(1, d-1)$ can be written in the form

$$f = x_1 g_1 + \ldots + x_n g_n,$$

where $g_i \in \mathbb{K}[u_1, \ldots, u_m]_{d-1}$. We say that $f$ is of *monomial square free type* if all $g_i$ are square free monomials. The associated algebra, $A = Q/\operatorname{Ann}_Q(f)$, is bigraded, has socle bidegree $(1, d-1)$ and we assume that $I_1 = 0$, so $\operatorname{codim} A = m + n$.

We enhance that the combinatoric structure inward bihomogeneous polynomials of monomial square free type, at least in low degree, allows us to give necessary and sufficient conditions in order to the associated algebra to be presented by quadrics. On the other hand, Gordan-Noether theory of forms with vanishing Hessian and the Lefschetz-Hessian criteria allow us to construct, in sufficiently large codimension, Artinian Gorenstein algebras presented by quadrics faling the WLP.

2.2. **The Hessian criteria for Strong and Weak Lefschetz properties.** Our point of view in order to study the Lefschetz properties for standard graded Artinian Gorenstein algebras, $A = Q/\operatorname{Ann}_Q(f)$, was inspired by [MW], where the authors proved a determinantal Hessian criterion, identifying Strong Lefschetz elements. In [Go] the first author studied and generalized class of examples given in [MW]. Now we prove a generalization of the Strong Lefschetz criterion using mixed Hessians. Let us introduce this natural object.

**Definition 2.8.** Let $A = Q/\operatorname{Ann}_Q(f) = \bigoplus_{k=0}^{d} A_k$ be a standard graded Artinian Gorenstein $\mathbb{K}$ algebra of socle degree $d$. Let $i \leq j \leq \frac{d}{2}$ be two integers and let $B_k = \{\alpha_1, \ldots, \alpha_s\}$ and $B_l = \{\beta_1, \ldots, \beta_t\}$ be bases of the $\mathbb{K}$-vector spaces $A_k$ and $A_l$ respectively. The (mixed) hessian matrix of $f$ of order $(k, l)$ is the matrix:

$$\operatorname{Hess}_f^{(k,l)} = (\alpha_i(\beta_j(f)))_{s \times t}.$$

We denote $\operatorname{Hess}_f^k := \operatorname{Hess}_f^{(k,k)}$, $\operatorname{hess}_f^k := \det(\operatorname{Hess}_f^k)$ and $\operatorname{hess}_f := \operatorname{hess}_f^1$.

**Remark 2.9.** We observe that, under the natural assumption that $\operatorname{Ann}_Q(f)_1 \neq 0$, the notation $\operatorname{hess}_f$ is consistent with the classical definition of Hessian, by taking $B_1 = \{X_1, \ldots, X_n\}$, the standard basis of the embedding.
If $A$ is bigraded, and if $B_k = \{\alpha_1, \ldots, \alpha_s\}$ and $B_l = \{\beta_1, \ldots, \beta_t\}$ are bases of the $\mathbb{K}$-vector spaces $A_{(k,l)}$ and $A_{(k',l')}$ respectively, we can also define $\operatorname{Hess}_f^{((k,l),(k',l'))} = (\alpha_i(\beta_j(f)))_{s \times t}$.

We now prove a generalization of [Wa1, Theorem 4] and [MW, Theorem 3.1].

**Theorem 2.10. (Hessian criteria for Strong and Weak Lefschetz elements)**
*Let $A = Q/\operatorname{Ann}_Q(f)$ be a standard graded Artinian Gorenstein algebra of codimension $r$ and socle degree $d$ and let $L = a_1 x_1 + \ldots + a_r x_r \in A_1$, such that $f(a_1, \ldots, a_r) \neq 0$. The map $\bullet L^{l-k} : A_k \to A_l$, for $k < l \leq \frac{d}{2}$, has maximal rank if and only if the (mixed) Hessian matrix $\operatorname{Hess}_f^{(k,d-l)}(a_1, \ldots, a_r)$ has maximal rank. In particular, we get the following:*

(1) **(Strong Lefschetz Hessian criterion,** [Wa1]**,** [MW]**)** *$L$ is a strong Lefschetz element of $A$ if and only if $\operatorname{hess}_f^k(a_1, \ldots, a_r) \neq 0$ for all $k = 1, \ldots, [d/2]$.*

(2) **(Weak Lefschetz Hessian criterion)** *$L \in A_1$ is a weak Lefschetz element of $A$ if and only if either $d = 2q+1$ is odd and $\mathrm{hess}_f^q(a_1, \ldots, a_r) \neq 0$ or $d = 2q$ is even and $\mathrm{Hess}_f^{(q-1,q)}(a_1, \ldots, a_r)$ has maximal rank.*

*Proof.* For finite dimensional $\mathbb{K}$-vector spaces $\mathbb{V}, \mathbb{W}$ we have the isomorphism:

$$\mathrm{Hom}_{\mathbb{K}}(\mathbb{V}, \mathbb{W}) \simeq \mathrm{Hom}_{\mathbb{K}}(\mathbb{V} \otimes \mathrm{Hom}_{\mathbb{K}}(\mathbb{W}, \mathbb{K}), \mathbb{K}).$$

Therefore, to any $\mathbb{K}$-linear map $T : \mathbb{V} \to \mathbb{W}$ we can associate, in a unique natural way, a bilinear form $B_T : \mathbb{V} \times \mathbb{W}^* \to \mathbb{K}$. Furthermore, $T$ has maximal rank if, and only if, $B_T$ has maximal rank. In our case of interest, the multiplication of the algebra induces the dual pairing, $A_l \times, A_{d-l} \to A_d \simeq \mathbb{K}$ via the identification $A_d \simeq \mathbb{K}$ given by $\omega \in A_d \leftrightarrow \omega(f) \in \mathbb{K}$. Hence, the linear map $\bullet L : A_k \to A_l$ induces a bilinear map $A_k \times A_{d-l} \to \mathbb{K}$ given by $(\alpha, \gamma) \mapsto \gamma.L^{l-k}.\alpha(f)$. If $\alpha_1, \ldots, \alpha_s \in A_k$ and $\gamma_1, \ldots, \gamma_t \in A_{d-l}$ are $\mathbb{K}$-linear bases, then the matrix of $B_T$ with respect to these bases is given by $[B_T] = (\gamma_i.L^{l-k}.\alpha_j(f))_{t\times s} = (L^{l-k}(\gamma_i\alpha_j(f)))_{t\times s}$. On the other side, $\mathrm{Hess}_f^{(k,d-l)} = [\gamma_j\alpha_i(f)]$, and we denote $g_{ij} := \gamma_j\alpha_i(f) \in R_{l-k}$. Since $L^{l-k}g_{ij} = (l-k)!g_{ij}(\underline{a})$, the result follows. The others claim follows from Remark 2.5. □

2.3. **Forms with vanishing Hessian.** Perazzo polynomials are the atoms of all known constructions of polynomials with vanishing hessian (see [Go] and [CRS] for more details). The following easy Proposition can be inferred from Gordan-Noether work. A proof of it can be found in [Ru, Section 7.2.1].

**Proposition 2.11. (Gordan-Noether criterion** [GN]**)** *Let $f \in \mathbb{K}[x_1, \ldots, x_n]$ be a homogeneous polynomial of degree $d$. Then $\mathrm{hess}_f = 0$ if and only if the partial derivatives of $f$ are algebraically dependent.*

**Definition 2.12.** $R = \mathbb{K}[x_1, \ldots, x_n, u_1, \ldots, u_m]$, for $m \geq 3$ and $n \geq 2$ be the polynomial ring and $Q$ be the associated ring of differential operators. A Perazzo polynomial of degree $d$ is a reduced polynomial $f \in R$ of the form

$$f = x_1g_1 + \ldots + x_ng_n + h \tag{1}$$

where $g_i \in \mathbb{K}[u_1, \ldots, u_m]_{d-1}$ for $i = 1, \ldots, n$ are algebraically dependent but linearly independent and $h \in \mathbb{K}[u_1, \ldots, u_m]_d$. The algebra $A = Q/\operatorname{Ann}_Q(f)$ is called a Perazzo algebra.

**Remark 2.13.** It is very easy to see that Perazzo polynomials have vanishing hessian, since the first derivatives $f_i = g_i$ for $i = 1, \ldots, n$ are algebraically dependent. Hence, by Gordan-Noether criterion, Proposition 2.11, $\mathrm{hess}_f = 0$. Notice that if $n > m$, then all polynomial of type 1 are Perazzo polynomials. By the Strong Lefschetz Hessian criterion, Theorem 2.10, Perazzo algebras do not have the SLP. Notice also that the construction begins in codimention five, this can be explained by Gordan-Noether Theorem, 2.14.

**Theorem 2.14.** [GN] *Let $X = V(f) \subset \mathbb{P}^N$ be a hypersurface, not a cone, given by a reduced polynomial $f$. If $N \leq 3$, then $\mathrm{hess}_f \neq 0$.*

**Corollary 2.15.** [MW] *Let $A$ be an standard graded Artinian Gorenstein algebra of socle degree $d \leq 4$ and codimension $r \leq 4$. Then $A$ has the SLP.*

## 3. Bigraded algebras, simplicial complexes and the main results

Let $f \in \mathbb{K}[x_1,\dots,x_n,u_1,\dots,u_m]_{(1,d-1)}$ be a bihomogeneous form of monomial square free type, $f = \sum_{i=1}^{n} x_i g_i$. Let us consider the homogeneous (or pure) simplicial complex $\mathcal{K} = \mathcal{K}(u)$ of dimension $d-1$ whose facets are given by the monomials $g_i$. The 0-skeleton will be referred as vertex set and we write $V = \{u_1,\dots,u_m\}$. We identify the 1-skeleton with a simple graph $\mathcal{K}_1 = (V,E)$, hence the 1-faces are called edges. Since $X_i(f) = g_i$, we identify each facet $g_i$ with the differential operator $X_i$. We denote by $e_k$ the number of $k$-faces, hence $e_0 = m$ and $e_{d-1} = n$. Let $A = Q/\operatorname{Ann}_Q(f)$ be the associated algebra, we suppose that $I_1 = 0$. We identify the faces of $\mathcal{K}$ with the dual differential operators, $u_i \leftrightarrow U_i$. If $p \in \mathbb{K}[u_1,\dots,u_m]$ is a square free monomial, we denote by $P \in \mathbb{K}[U_1,\dots,U_m]$ the dual differential operator $P = p(U_1,\dots,U_m)$, notice that $P(p) = 1$.

**Theorem 3.1.** *Let $f = \sum_{i=1}^{n} x_i g_i \in \mathbb{K}[x_1,\dots,x_n,u_1,\dots,u_m]_{(1,d-1)}$ be a bihomogeneous polynomial of monomial square-free type, let $\mathcal{K}$ be the associated simplicial complex and let $A = Q/\operatorname{Ann}_Q(f)$ be the standard bigraded Artinian Gorenstein algebra of codimension $m+n$ and socle bidegree $(1,d-1)$. Then $A = \bigoplus_{k=0}^{d} A_k$ where $A_k = A_{(0,k)} \oplus A_{(1,k-1)}$. Moreover, $A_{(0,k)}$ has a basis identified with the $k$ faces of $\mathcal{K}$, hence $\dim A_{(0,k)} = e_k$. By duality, $A^*_{(1,k-1)} \simeq A_{(0,d-k)}$, and a basis for $A_{(1,k-1)}$ can be chosen by taking, for each $d-k$-face of $\mathcal{K}$, a monomial $X_i\tilde{G}_i$ such that $X_i\tilde{G}_i(f)$ represents this $d-k$-face. In particular, the Hilbert vector of $A$ is given by $h_k = \dim A_k = e_k + e_{d-k}$. Furthermore, $I = \operatorname{Ann}_Q(f)$ is a binomial ideal generated by $(X_1,\dots,X_n)^2$, all the monomials in $U_i$ that do not represent faces of $\mathcal{K}$, all monomials $X_iF_i$ where $f_i$ does not represent a subface of $g_i$ and all binomials $X_i\tilde{G}_i - X_j\tilde{G}_j$ where $g_i = \tilde{g}_i g_{ij}$ and $g_j = \tilde{g}_j g_{ij}$ and $g_{ij}$ represents a common subface of $g_i, g_j$.*

*Proof.* It is easy to see that $A_{(0,k)}$ is generated by the monomials that represent $k$-faces, since they are the only ones that do not annihilate $f$. Now we show that they are linearly independent over $\mathbb{K}$. For any $k$-face $\omega$, let $\Omega$ be the monomial of $Q_{(0,k)}$, let $\Omega_1,\dots,\Omega_s$ be all of them. Since $\Omega(f) = \sum_{i=1}^{n} x_i\Omega(g_i)$, if we take any linear combination

$$0 = \sum_{j=1}^{s} c_j\Omega_j(f) = \sum_{j=1}^{s} c_j \sum_{i=1}^{n} x_i\Omega_j(g_i) = \sum_{i=1}^{n} x_i \sum_{j=1}^{s} c_j\Omega_j(g_i).$$

We get $\sum_{j=1}^{s} c_j\Omega_j(g_i) = 0$ for all $i = 1,\dots,n$. Fixed $i$, $\Omega_j(g_i)$ are distinct monomials or zero, but for each $j$ there is a $i$ such that $\Omega_j(g_i) \neq 0$, therefore $c_j = 0$ for all $j = 1,\dots,s$. The other assertions concerning $A$ are now clear.
Notice that $I_{(i,j)} = Q_{(i,j)}$ for all $i \geq 2$ and it is generated by $I_{(2.0)} = (X_1,\dots,X_n)^2$. Now we

describe $I_{(0,k)}$ and $I_{(1,k-1)}$. Consider the sequence giving by evaluation:

$$0 \to I_{(0,k)} \to Q_{(0,k)} \to A_{(1,d-1-k)} \to 0.$$

Since $\dim A_{(1,d-1-k)} = \dim A^*_{(1,d-1-k)} = \dim A_{(0,k)} = e_k$, we get $\dim I_{(0,k)} = \dim Q_{(0,k)} - e_k$. Since $\dim A_{(0,k)} = e_k$ and it has a basis given by the $k$-faces of $\mathcal{K}$ and since all the other $\dim Q_{(0,k)} - e_k$ monomials are linearly independent elements of $I_{(0,k)}$, they form a basis for it. Consider the sequence giving by evaluation:

$$0 \to I_{(1,k-1)} \to Q_{(1,k-1)} \to A_{(0,d-k)} \to 0.$$

We have $\dim I_{(1,k-1)} = Q_{(1,k-1)} - e_{d-k}$. Let us write $Q_{(1,k-1)} = \tilde{I}_{(1,k-1)} \oplus \tilde{Q}_{(1,k-1)}$ where $\bar{I}_{(1,k-1)}$ is the $\mathbb{K}$-vector space spanned by the monomials $X_iF_i$ where $F_i$ does not represent a subface of $G_i$. Of course $\bar{I}_{(1,k-1)} \subset I_{(1,k-1)}$ and $\tilde{Q}_{(1,k-1)}$ is spanned by all the monomials $X_i\tilde{G}_i$ where $\tilde{G}_i$ is a subface of $G_i$. The exact sequence given by evaluation restricted to $\tilde{Q}_{(1,k-1)}$ becomes

$$0 \to \tilde{I}_{(1,k-1)} \to \tilde{Q}_{(1,k-1)} \to A_{(0,d-k)} \to 0.$$

Hence, $I_{(1,k-1)} = \tilde{I}_{(1,k-1)} \oplus \bar{I}_{(1,k-1)}$, since $X_i\tilde{G}_i(f)$ is a face of $\mathcal{K}$, $\tilde{I}_{(1,k-1)}$ is generated by the binomials $X_i\tilde{G}_i - X_j\tilde{G}_j$ such that $X_i\tilde{G}_i(f) = g_{ij} = X_j\tilde{G}_j(f)$ where $g_{ij}$ is a common subface of $g_i, g_j$, $g_i = \tilde{g}_i g_{ij}$ and $g_j = \tilde{g}_j g_{ij}$. The result follows. □

**Definition 3.2.** Let $\mathcal{K}$ be a homogeneous simplicial complex of dimension $d-1$. We say that $\mathcal{K}$ is facet connected if for any pair of facets $F, F'$ of $\mathcal{K}$ there exists a sequence of facets, $F_0 = F, F_1, \ldots, F_s = F'$ such that $F_i \cap F_{i+1}$ is a $(d-2)$-face. We say that $\mathcal{K}$ is upper closed if for all complete subgraphs $H = K_l \subset \mathcal{K}_1$ there is a $l$-face $F \in \mathcal{K}_l$ such that $H$ is the first skeleton of $F$. In particular $\mathcal{K}_1$ does not contain any $K_d$.

**Theorem 3.3.** *Let $f \in \mathbb{K}[x_1, \ldots, x_n, u_1, \ldots, u_m]_{(1,d-1)}$ be a bihomogeneous polynomial of monomial square-free type, let $\mathcal{K}$ be the associated simplicial complex and let $A = Q/\operatorname{Ann}_Q(f)$ be the standard bigraded Artinian Gorenstein algebra. $A$ is presented by quadrics if and only if $\mathcal{K}$ is facet connected and upper closed.*

*Proof.* Suppose that $\mathcal{K}$ is upper closed and facet connected. By Theorem 3.1, $I = \operatorname{Ann}_Q(f)$ it is enough to consider the monomials in the $U_i$ that does not represent a face of $\mathcal{K}$, monomials $X_iF_i$ where $F_i$ is a monomial in the $U_j$ that does not represent a subface of $G_i$ and the binomials $X_i\tilde{G}_i - X_j\tilde{G}_j$ where $X_i\tilde{G}_i(f) = g_{ij} = X_j\tilde{G}_j(f)$ is a common subface of $g_i, g_j$. Let $M = U_1^{e_1} \ldots U_m^{e_m}$ be a monomial such that $M(f) = 0$, if $e_i \geq 0$ for some $i = 1, \ldots, m$, then $M = U_i^2\tilde{M}$ and since $U_i^2 \in I_2$, $M \in I_2Q$, hence we can consider $M$ square free and it does not represents a face of $\mathcal{K}$. In this case, $M$ represents a $K_l$ with the same vertex set of $G$, since, by hypothesis, for $3 \leq l \leq d-2$ all $K_l \subset G$ comes from a $l$-face of $\mathcal{K}$ and since $G$ does not contain a $K_{d-1}$ as subgraph, there exists $U_iU_j$ in $M$ such that $U_iU_j(f) = 0$ and $M = U_iU_j\tilde{M} \in I_2Q$.

Let $\Omega = X_iM$ with $M = U_1^{e_1} \ldots U_m^{e_m}$ be a monomial such that $\Omega(f) = M(g_i) = 0$, if $e_i \geq 2$ for some $i = 1, \ldots, m$, then $\Omega = U_i^2\tilde{\Omega} \in I_2Q$, so we can suppose that $M$ is square free and it does not represent a subface of $g_i$, hence there is a $U_j$ in $M$ that does not belongs to $G_i$, yielding $\Omega = X_iU_j\tilde{M} \in I_2Q$.

To finish the proof, consider the binomials $X_i\tilde{G}_i - X_j\tilde{G}_j$ where $X_i\tilde{G}_i(f) = g_{ij} = X_j\tilde{G}_j(f)$ and

$g_{ij}$ is a common subface of $g_i, g_j$. If $\tilde{G}_i$ and $\tilde{G}_j$ are subfaces of the facets $G_i, G_j$ respectively and if $g_{ij} \subset G_i \cap G_j$ and the intersection is a $d-2$-face, then there are only two vertices they do not share, say $u_i, u_j$, $\tilde{G}_i = U_i G_{ij}$ and $\tilde{G}_j = U_j G_{ij}$ and finally $X_i\tilde{G}_i - X_j\tilde{G}_j = (X_iU_i - X_jU_j)G_{ij} \in I_2Q$. In the general case, by the facet connection of $\mathcal{K}$, there exists a sequence of facets $G_{i_0} = G_i, G_{i_1}, \ldots, G_{i_s} = G_j$ such that the intersection of two consecutive facets is a $d-2$-face, hence $X_i\tilde{G}_i - X_{i_1}\tilde{G}_{i_1}, X_{i_1}\tilde{G}_{i_1} - X_{i_2}\tilde{G}_{i_2}, \ldots, X_{i_s}\tilde{G}_{i_s} - X_j\tilde{G}_j \in I_2Q$, summing up we get the desired result.

Conversely, if $\mathcal{K}$ is not facet connected, let $g_j, g_j$ be two facets that can not be facet connected and let $g_{ij} = \gcd(g_i g_j)$. By Theorem 3.1 it is easy to see that $X_i\tilde{G}_i - X_j\tilde{G}_j$ is a minimal generator of $I$ where $g_i = \tilde{g}_i g_{ij}$ and $g_j = \tilde{g}_j g_{ij}$. If $\mathcal{K}$ is not upper closed, then there is a complete subgraph $K_s \subset G$ that does not came from a $s$-face of $\mathcal{K}$. In this case, if we choose $s$ to be minimal, then by Theorem 3.1 the monomial $M = \prod_{v \in V(K_s)} v$ is a minimal generator of $I$.

□

We introduce the following complexes inspired by the famous Turan Theorem characterizing maximal graphs not containing a $K_{d-1}$ as the $(d-1)$-partite complete graph $K(a_1, \ldots, a_{d-1})$ whith $|a_i - a_j| \leq 1$. Here is the definition.

**Definition 3.4.** Let $2 \leq a_1 \leq \ldots \leq a_{d-1}$ be integers, the Turan complex of order $a_1, \ldots, a_{d-1}$, $\mathcal{K} = \mathcal{TK}(a_1, \ldots, a_{d-1})$, is the Homogeneous simplicial complex whose facets set is the cartesian product $\pi = \prod_{i=1}^{d-1} \{1, 2, \ldots, a_i\}$. The Turan polynomial of order $a_1, \ldots, a_{d-1}$ it the multihomogeneous polynomial $f = f_{\mathcal{K}} = \sum_{\alpha \in \pi} x_\alpha u_\alpha \in R = \mathbb{K}[x_\alpha, u_{(i,j_i)}]_{\alpha \in \pi, 1 \leq i \leq d-1, 1 \leq j_i \leq a_i}$, where $\alpha = (j_1, \ldots, j_{d-1}) \in \pi$ and $u_\alpha = u_{(1,j_1)} \ldots u_{(d-1,j_{d-1})}$. The Turan algebra of order $(a_1, \ldots, a_{d-1})$ is $TA(a_1, \ldots, a_{d-1}) = A_{\mathcal{K}} = Q/\operatorname{Ann}_Q(f)$.

**Remark 3.5.** Notice that the number of $k$-faces of a Turan complex is $e_k = s_k$ where $s_k = s_k(a_1, \ldots, a_{d-1})$ is the symmetric function of order $k$. By Theorem 3.3, the Hilbert vector of the Turan algebra $TA(a_1, \ldots, a_{d-1})$ is given by $h_k = s_k + s_{d-k}$. It is easy to verify that Turan complexes are facet connected and upper closed.

**Lemma 3.6.** *Let $f \in R_{(1,d-1)}$ be a bihomogeneous polynomial of monomial square free type, let $\mathcal{K}$ be the associated simplicial complex and $A$ the associated algebra. Let $e_k$ the number of $k$-faces of $\mathcal{K}$. Then the map $\bullet L : A_{k-1} \to A_k$, for $k \leq \frac{d}{2}$, is injective for a general $L \in A_1$ if, and only if* $\operatorname{rk} \operatorname{Hess}_f^{((1,k-2),(0,d-k))} = e_{d-k+1}$ *and* $\operatorname{rk} \operatorname{Hess}_f^{((0,k-1),(1,d-k-1))} = e_{k-1}$.

*Proof.* Since $A_k = A_{(1,k-1)} \oplus A_{(0,k)}$, and since by Theorem 3.3 $\dim A_{(0,k)} = e_k$ and $\dim A_{(1,k-1)} = \dim A_{(0,d-k)} = e_{d-k}$, with a choice of bases consistent with the decomposition as direct sum, we have:

$$\operatorname{Hess}_f^{(k-1,d-k)} = \begin{bmatrix} 0 & \operatorname{Hess}_f^{((1,k-2),(0,d-k))} \\ \operatorname{Hess}_f^{((0,k-1),(1,d-k-1))} & \operatorname{Hess}_f^{((0,k-1),(0,d-k))} \end{bmatrix}_{(e_{d-k+1}+e_{k-1}) \times (e_k + e_{d-k})},$$

where the matrices $\mathrm{Hess}_f^{((1,k-2),(0,d-k))}$ and $\mathrm{Hess}_f^{((0,k-1),(1,d-k-1))}$ have order $e_{d-k+1} \times e_{d-k}$ and $e_{k-1} \times e_k$ respectively.

The injectivity of $\bullet L : A_{k-1} \to A_k$ implies $e_{d-k+1}+e_{k-1} \leq e_{d-k}+e_k$ and $\operatorname{rk}\mathrm{Hess}_f^{(k-1,d-k)} = e_{d-k+1} + e_{k-1}$. By the shape of the matrix this maximal rank can be achieved if and only if $\operatorname{rk}\mathrm{Hess}_f^{((1,k-2),(0,d-k))} = e_{d-k+1}$ and $\operatorname{rk}\mathrm{Hess}_f^{((0,k-1),(1,d-k-1))} = e_{k-1}$.

Conversely, if $\operatorname{rk}\mathrm{Hess}_f^{((1,k-2),(0,d-k))} = e_{d-k+1}$ and $\operatorname{rk}\mathrm{Hess}_f^{((0,k-1),(1,d-k-1))} = e_{k-1}$, then $\operatorname{rk}\mathrm{Hess}_f^{(k-1,d-k)} = e_{d-k+1} + e_{k-1}$ yielding the desired result. □

**Lemma 3.7.** *Let $d \geq 3$ be an integer and consider the Turan complex $\mathcal{TK}(2^{(d-1)}) := \mathcal{TK}(2,\ldots,2)$ of dimension $d-1$. Let $f$ be the associated form. Then* $\operatorname{rk}\mathrm{Hess}_f^{((1,0),(0,d-2))} < 2^{d-1}$.

*Proof.* Let us write $\mathcal{K} = \mathcal{TK}(2^{(d-1)})$. First of all note that the rows of $\mathrm{Hess}_f^{((1,0),(0,d-2))}$ are indexed by the $2^{d-1}$ facets $x_\alpha$ of $\mathcal{K}$ and the columns are indexed by the $(d-2)$-faces $F$ of $\mathcal{K}$. A non zero element of $\mathrm{Hess}_f^{((1,0),(0,d-2))}$ is a degree one monomial representing the remaining vertex of the facet $u_\alpha$ that does not belongs to the $(d-2)$-face $F$. For instance, every column $F$ has only two non zero elements, say $u_{ij}$ and $u_{kj}$ representing the remaining vertex of the two faces that contain $F$. Furthermore, the other non zero elements of the rows $i$ and $k$ are the same.

If we multiply every row indexed by $x_\alpha$, with $\alpha = (j_1,\ldots,j_{d-1})$ where $j_i \in \{0,1\}$ by $(-1)^{j+1+\ldots+j_{d-1}}$, and by all the variables that do not figure in the row. Then we get a matrix $M$ such that every column $j$ has only two non zero elements and they are opposite, say $M_j$ and $-M_j$. Summing up the rows, the result follows. □

**Lemma 3.8.** *Let $\mathcal{K}$ be a pure simplicial complex of dimension $d-1$, let $n$ be the number of facets of $\mathcal{K}$, let $f$ be the associated bigraded polynomial of monomial square free type and let $A$ be the associated standard bigraded algebra of socle bidegree $(1,d-1)$. Let $v \in V(\mathcal{K})$ be a vertex and denote $\mathcal{K}' = \mathcal{K} \setminus v$ be the complex obtained from $\mathcal{K}$ by deleting $v$, let $n'$ be the number of facets of $\mathcal{K}'$, let $f'$ be the associated polynomial and $A'$ the associated algebra. Then*

$$\operatorname{rk}\mathrm{Hess}_f^{((1,0),(0,d-2))} = n \Rightarrow \operatorname{rk}\mathrm{Hess}_{f'}^{((1,0),(0,d-2))} = n'.$$

*Proof.* Let us choose an ordered basis of $A_{(0,1)}$ such that the last $n'$ vectors represent the faces containing $v$. Let us choose a basis of $A_{(0,d-2)}$ in such a way that the first vectors represent $d-2$ faces that does not contain $v$ and the last vectors the faces that contain $v$. The Matrix $\mathrm{Hess}_f^{((1,0),(0,d-2))}$ with respect to this basis is

$$\mathrm{Hess}_f^{((1,0),(0,d-2))} = \begin{bmatrix} * & * \\ 0 & \mathrm{Hess}_{f'}^{((1,0),(0,d-2))} \end{bmatrix}.$$

The zero sub-matrix occurs by our choice of ordered basis. In fact, if $X_i$ represent a face not containing $v$, then $X_i(f)$ does not contain the variable $v$ and since the first vector of $A_{(0,d-2)}$ contain $v$, the derivative is zero. The result easily follows. □

We now present families of counterexamples to Migliore-Nagel injective conjecture that occur in large codimension with respect to the socle degree. It is surprising that if the codimension is very large, the Hilbert vector becomes non unimodal in the first step.

**Corollary 3.9.** *Let $A = TA(a_1, \ldots, a_{d-1})$ be the Turan algebra of order $(a_1, \ldots, a_{d-1})$ with $2 \leq a_1 \leq a_2 \leq \ldots \leq a_{d-1}$. Then $A$ is presented by quadrics and for all $L \in A_1$ the map $\bullet L : A_1 \to A_2$ is not injective. Furthermore, if $a_1 \approx \ldots \approx a_{d-1}$ are large enough, then* $\mathrm{Hilb}(A)$ *is not unimodal in the first step, that is,* $\dim A_1 > \dim A_2$.

*Proof.* First of all it is easy to see that $\mathcal{TK}(a_1, \ldots, a_{d-1})$ is facet connected and upper closed, hence, by Theorem 3.3, $TA(a_1, \ldots, a_{d-1})$ is presented by quadrics. By Remark 3.5 we have $e_k = s_k$, the symmetric functions. For $a_1 = \ldots = a_{d-1} = 2$, the result follows by Lemma 3.7 and Lemma 3.6, for each $a_i > 2$ we can delete vertex until obtain $\mathcal{TK}(2^{d-1})$ and by Lemma 3.8 and Lemma 3.6 the result follows.

If $a_1 \approx \ldots \approx a_{d-1} \approx a$ are large enough, then, by a trivial Calculus I argument:

$$\dim A_1 \approx (d-1)a + a^{d-1} > \binom{d-1}{2} a^2 + (d-1)a^{d-2} \approx \dim A_2.$$

In this case, the Hilbert vector $\mathrm{Hilb}(A)$ is not unimodal in the first step. □

## 4. Indutive constructions and bounds for the WLP conjecture

By Corollary 3.9, for each socle degree $d \geq 3$ and for large codimension there exist Artinian Gorentein algebras such that for all $L \in A_1$, the map $\bullet : A_1 \to A_2$ is not injective, of course they are counterexamples for both MN- conjectures. In this section we find lower bounds for the codimension, as a function of the socle degree, for the existence of counterexamples for the WLP conjecture.

### 4.1. **An indutive construction.**

**Lemma 4.1.** *Let $f \in R = \mathbb{K}[x_1, \ldots, x_r]$ be a homogeneous polynomial of degree $d$ and let $\tilde{f} = uf \in \tilde{R} = \mathbb{K}[x_1, \ldots, x_r, u]$. Let $Q$ and $\tilde{Q}$ be the rings of differential operators associated to $R$ and $\tilde{R}$ respectively. Then*

$$\mathrm{Ann}_{\tilde{Q}}(\tilde{f}) = \mathrm{Ann}_Q(f)\tilde{Q} + U^2\tilde{Q} \subset \tilde{Q}.$$

*In particular, if $A = Q/\mathrm{Ann}_Q(f)$ is presented by quadrics, then so it is for $\tilde{A} = \tilde{Q}/\mathrm{Ann}_{\tilde{Q}}(\tilde{f})$.*

*Proof.* It is easy to see that if $\alpha \in \mathrm{Ann}_Q(f)$, then $\alpha \in \mathrm{Ann}_{\tilde{Q}}(\tilde{f})$, and also $U^2 \in \mathrm{Ann}_{\tilde{Q}}(\tilde{f})$, hence $I = \mathrm{Ann}_Q(f)\tilde{Q} + U^2\tilde{Q} \subset \mathrm{Ann}_{\tilde{Q}}(\tilde{f})$. To prove the equality, let $\overline{\alpha} \in \mathrm{Ann}_{\tilde{Q}}(\tilde{f})/I$, then we can write:

$$\overline{\alpha} = \overline{\beta} + U\overline{\gamma}.$$

Where $\beta, \gamma \in Q/I \subset \tilde{Q}/I$. Therefore:

$$\alpha(\tilde{f}) = \beta(f) + U\gamma(f) = 0.$$

Which give us $\overline{\beta} = \overline{\gamma} = 0$, hence $\overline{\alpha} = 0$ and the result follows. □

**Lemma 4.2.** *Let $f \in R = \mathbb{K}[x_1, \dots, x_r]$ be a homogeneous polynomial of degree $d$ and let $\tilde{f} = uf \in \tilde{R} = \mathbb{K}[x_1, \dots, x_r, u]$. Let $Q$ and $\tilde{Q}$ be the ring of differential operators associated to $R$ and $\tilde{R}$ respectively. Then we have the following decomposition as $\mathbb{K}$ vector spaces:*

$$\tilde{A}_k = A_k \oplus A_{k-1}U.$$

*Proof.* Let $\{\beta_1, \dots, \beta_s\} \subset A_k$ be a $\mathbb{K}$-basis of $A_k$ and let $\{\gamma_1, \dots, \gamma_l\} \subset A_{k-1}$ be a $\mathbb{K}$-basis of $A_{k-1}$. We claim that $\{\beta_1, \dots, \beta_s, U\gamma_1, \dots, U\gamma_l\} \subset \tilde{A}_k$ is a $\mathbb{K}$-basis of $\tilde{A}_k$.

(i) Linear independence. Suppose that

$$b_1\beta_1 + \ldots + b_s\beta_s + c_1U\gamma_1 + \ldots + c_lU\gamma_l = 0.$$

Hence, by Lemma 4.1 $b_1\beta_1 + \ldots + b_s\beta_s = 0$ implying $b_1 = \ldots = b_s = 0$, in the same way $c_1U\gamma_1 + \ldots + c_lU\gamma_l = 0$ implying $c_1 = \ldots = c_l = 0$.

(ii) Spanning. Let $\alpha \in \tilde{A}_k$, by Lemma 4.1, $\alpha = \beta + U\gamma$, with $\beta \in A_k$ and $\gamma \in A_{k-1}$. Therefore $\beta = b_1\beta_1 + \ldots + b_s\beta_s$ and $\gamma = c_1U\gamma_1 + \ldots + c_lU\gamma_l$ since $\{\beta_1, \dots, \beta_s\}$ is a $\mathbb{K}$-basis of $A_k$ and $\{\gamma_1, \dots, \gamma_l\}$ is a $\mathbb{K}$-basis of $A_{k-1}$.

□

**Definition 4.3.** Let $h = (1, h_1, h_2, \dots, h_{d-1}, 1)$ be a Gorenstein Hilbert vector, the previous Lemma 4.2 allows to define the Hilbert vector $\sigma(h) := (1, 1 + h_1, h_1 + h_2, \dots, h_{d-1} + 1, 1)$. Iterating Lemma 4.2 $k$ times, we define $\sigma^{(k)}$.

**Corollary 4.4.** *There exist standard bigraded Artinian Gorenstein algebras $A$, presented by quadrics, of socle bidegree $(1, d-1)$, codimension $r = d$, $r = d+2$ or $r \geq d$ and for all $d \geq 3$. Furthermore, the Hilbert vector of $A$ can be choosen to be* $\mathrm{Hilb}(A) = \sigma^{(d-3)}(1, r, r, 1)$.

*Proof.* By Theorem 3.3 the result follows for $d = 3$. Indeed, for odd $r \geq 3$ it is enough to take any tree, for even $r \geq 8$ any circuit. By induction on $d \geq 3$ and by Lemma 4.2 and Lemma 4.1 the result follows. □

The following Lemma will be useful in the next sections. If $\mathcal{K}$ is a simplicial complex and $l \in \mathcal{K}_{d-2}$ is a $d-2$-face, we construct the complex $\mathcal{K}'$ attaching a new facet from $l$ and a new vertex. This facet is said to be a *leaf*. Notice that $e'_0 = e_0 + 1$, $e'_i - e_i \geq 2$ for $2 \leq i \leq d-2$ and $e'_d = e_d + 1$.

**Lemma 4.5.** *Let $\mathcal{K}$ be a $d-1 \geq 2$ dimensional simplicial complex that represents an algebra $A$, presented by quadrics such that $d_1 \leq d_2$ and $e_{d-1} \leq e_{d-2}$. Let $\mathcal{K}'$ be the simplicial complex constructed from $\mathcal{K}$ attaching a leaf. Then the algebra $A'$ given by $\mathcal{K}'$ is presented by quadrics. Furhtermore, $A$ satisfies the injective conjecture if, and only if $A'$ also satisfies.*

*Proof.* It is easy to see that $A'$ is presented by quadrics by Theorem 3.3. By Lemma 3.6, since $e_1 \leq e_2$ and $e_{d-1} \leq e_{d-2}$, we have that $A$ safisfies the injective conjecture if, and only if $\operatorname{rk} \mathrm{Hess}_f^{((1,0),(0,d-2))} = e_{d-1}$ and $\operatorname{rk} \mathrm{Hess}_f^{((0,1),(1,d-3))} = e_1$. Since attaching a leaf we still have $e'_1 \leq e'_2$ and $e'_{d-1} \leq e'_{d-2}$ and since it does not alters the fact of the rank of the desired mixed Hessians to be maximal, the result follows. □

4.2. **Artinian Gorenstein algebras with odd socle degree.** Let $A$ be a standard graded Artinian Gorenstein algebra with socle degree three, then $A = Q/\operatorname{Ann}_Q(f)$ with $f \in R$ a homogeneous polynomial of degree 3. Theorem 2.10 applied to this case tells us that $A$ satisfies the WLP if and only if $\operatorname{hess}_f \neq 0$.

By a result due to Dimca-Papadima, see see [DP, Thm. 1], if $f$ is not a reduced polynomial and $\tilde{f}$ is its radical, then $\operatorname{hess}_f = 0$ if and only if $\operatorname{hess}_{\tilde{f}} = 0$. For quadratic polynomials not a cone, $\operatorname{hess}_{\tilde{f}} \neq 0$, so we can restrict ourselves to reduced cubic polynomials. Furthermore, if $f = f_1 f_2$ and $\operatorname{hess}_f = 0$, then all the components of $X = V(f) \subset \mathbb{P}^n$ are developable, yielding $\operatorname{hess}_{f_i} \equiv 0 \pmod{f_i}$, in this case $f = l_1 l_2 l_3$ and $X$ is an arrangement of hyperplanes passing through a $\mathbb{P}^{N-2}$, which is a cone as soon as $N \geq 2$. So, from now on we can restric ourselves to the case that $f$ is an absolutely irreducible polynomial.

**Theorem 4.6.** [GRu] *Perazzo hypersurfaces are not cones and have vanishing Hessian. Suppose that $N \leq 6$, and let $X = V(f) \subset \mathbb{P}^N$ be an irreducible cubic hypersurface which is not a cone and such that* $\operatorname{hess}_f = 0$. *Then, up to a projective transformation, $f$ is a Perazzo polynomial.*

**Corollary 4.7.** *Let $A$ be a standard graded Artinian Gorenstein $\mathbb{K}$-algebra of socle degree* 3. *If $A$ is presented by quadrics and* $\operatorname{codim} A \leq 7$, *then $A$ satisfy the WLP.*

*Proof.* Suppose that $A$ does not have the $WLP$. Then by the Hessian criterion, Theorem 2.10, $\operatorname{hess}_f = 0$. For $N \leq 6$, by Theorem 4.6 and by [GRu, Thm. 5.2,5.3,5.4], if $f$ is an irreducible cubic polynomial such that $\operatorname{hess}_f = 0$, then, up to a projective transformation, either $f = x_0x_3^2 + x_1x_3x_4 + x_2x_4^2 + h$ with $h \in \mathbb{K}[x_3, \ldots, x_N]$ or $N = 6$ and $f = x_0g_0 + \ldots + x_3g_3 + h$ with $h \in \mathbb{K}[x_4, x_5, x_6]$. Let us suppose, without loss of generality, that $x_4^2$ occurs as monomial only in $g_2$. In both cases, if we consider the associated standard graded Artinian Gorenstein one can verify directly that $X_2^3 \in \operatorname{Ann}_Q(f)$ is a minimal generator. □

The next example was treated from the geometric point of view in [GRu, p. 803, Example 6]. It is the minimal counter-example for the MN-conjecture.

**Example 4.8.** In $\mathbb{P}^7$ consider the cubic hypersurface $X = V(f) \subset \mathbb{P}^7$, given by

$$f = \begin{vmatrix} x_0 & x_1 & x_2 \\ x_3 & x_4 & x_5 \\ x_6 & x_7 & 0 \end{vmatrix} \in \mathbb{K}[x_0, \ldots, x_7].$$

As pointed out in [GRu, p. 803, Example 6], $X$ represents a tangent section of the secant variety of the Segre variety $\operatorname{Seg}(\mathbb{P}^2, \mathbb{P}^2) \subset \mathbb{P}^8$. After a linear change of coordinates we can rewrite $f$ as a Perazzo polynomial of monomial square free type:

$$f = x_1u_1u_2 + x_2u_2u_3 + x_3u_3u_4 + x_4u_4u_1 \in R = \mathbb{K}[x_1, x_2, x_3, x_4, u_1, u_2, u_3, u_4].$$

Notice that $f_1f_3 = u_1u_2u_3u_4 = f_2f_4$, hence by the Gordan-Noether criterion, Proposition 2.11, $\operatorname{hess}_f = 0$. Let $A = Q/\operatorname{Ann}_Q(f)$ be the associated algebra, of codimension $\operatorname{codim} A = 8$ and socle degree 3. By the Hessian Lefschetz criterion, Theorem 2.10, $A$ does not has the WLP. On the other side, since it graph is a square, by Theorem 3.3 it is presented by quadrics. Indeed, one can verify that

$$I = (u_4^2, u_2u_4, x_2u_4, x_1u_4, u_3^2, u_1u_3, x_4u_3, x_1u_3, u_2^2, x_4u_2, x_3u_2, x_2u_2 - x_3u_4, x_1u_2 - x_4u_4,$$
$$u_1^2, x_4u_1 - x_3u_3, x_3u_1, x_2u_1, x_1u_1 - x_2u_3, x_4^2, x_3x_4, x_2x_4, x_1x_4, x_3^2, x_2x_3, x_1x_3, x_2^2, x_1x_2, x_1^2).$$

**Example 4.9.** Consider the algebras $A = Q/\operatorname{Ann}_Q(f)$ of codimension $r = 9, 11$. For $r = 9$, take $f = x_1u_1u_2 + x_2u_2u_3 + x_3u_3u_4 + x_4u_4u_1 + w^2u_1$ and for $r = 11$, take $f = x_1u_1u_2 + x_2u_2u_3 + x_3u_3u_4 + x_4u_4u_1 + x_5u_5u_1 + w^2u_1$. For both we have, $f_1f_3 = f_2f_4$, hence by Gordan-Noether criterion and the Hessian criterion, $A$ does not have the WLP. One can verify that in all the cases $\operatorname{Ann}_Q(f)$ is generated by quadrics.

**Lemma 4.10.** *Let $R = \mathbb{K}[u_1, \dots, u_m]$ be the polynomial ring. Let $G = (V, E)$ be a connected graph such that $V = \{u_1, \dots, u_m\}$ and $E$ is given by square free quadratic monomials. If $|E| = |V|$, then $G$ has a unique circuit $C$ and furthermore:*

1. *If $|C|$ is even, then $\det \nabla G = 0$;*
2. *If $|C|$ is odd, then $\det \nabla G \neq 0$*

*Proof.* First of all let us show that the gradient matrix of any tree has maximal rank. Let $T = (V', E')$ be a tree where $V' = \{u_1, \dots, u_m\}$ and $E' = \{g_1, \dots, g_{m-1}\}$. By induction on $m \geq 2$, the result is trivial for $m = 2$. Let us suppose that for any tree with $|V'| = m \geq 2$, the gradient matrix has maximal rank. Let $\tilde{T}$ be a tree with $|\tilde{T}| = m + 1$, $\tilde{T} = T \cup g_m$ where $g_m = u_j u_{m+1}$, hence

$$\nabla \tilde{T} = \begin{bmatrix} \nabla T & * \\ 0 & u_j \end{bmatrix}.$$

And the claim follows.
Let $T \subset G$ be a generating tree of $G$, then $T = (V, E')$ with $|E'| = |V| - 1$, since $|V| = |E|$, $G$ contains a unique circuit, say $C = \{u_1u_2, u_2u_3, \dots, u_{k-1}u_k, u_ku_1\}$ and let us suppose that $E = E' \cup u_ku_1$.

Since $G \setminus u_1u_k = T$ is a tree, $\nabla G = (\nabla T | \nabla g_m)$ where

$$\nabla T = \begin{bmatrix} u_2 & 0 & \dots & 0 & * \\ u_1 & u_3 & \dots & 0 & * \\ 0 & u_2 & \dots & 0 & * \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & u_n & * \\ 0 & 0 & \dots & u_{n-1} & * \\ 0 & 0 & \dots & 0 & N \end{bmatrix} \text{ and } \nabla G = \begin{bmatrix} u_2 & 0 & \dots & 0 & * & u_k \\ u_1 & u_3 & \dots & 0 & * & 0 \\ 0 & u_2 & \dots & 0 & * & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & \dots & u_n & * & 0 \\ 0 & 0 & \dots & u_{n-1} & * & u_1 \\ 0 & 0 & \dots & 0 & N & 0 \end{bmatrix}.$$

Where in the last row 0 represents $0_{(m-k,1)}$ and $N$ is a square matrix of order $m - k$ such that $\det(N) \neq 0$, since $\nabla T$ has maximal rank. Using the Laplace expansion, we get

$$\det(\nabla G) = \det(N) u_1u_2 \dots u_n(1 + (-1)^{k-1}).$$

The result follows. □

**Theorem 4.11.** *Let $f \in \mathbb{K}[x_1, \dots, x_n, u_1, \dots, u_m]_{(1,2)}$ be a bigraded cubic polynomial of monomial square free type and let $G = (E, V)$ be the associated graph. Then $A$ is presented by quadrics if, and only if, $G$ is connected and triangle free, in this case we have the following possibilities:*

1. *If $G$ is a tree, then $A$ has the WLP;*
2. *If $G$ contains only one circuit $|C|$, then either*
   (a) *$|C|$ is even and $A$ does not have the WLP or*
   (b) *$|C| \geq 5$ is odd and $A$ has the WLP*

(3) *If $G$ contains at least two circuits, then $A$ does not have the WLP.*

*Proof.* The characterization of the graphs that represent algebras presented by quadrics follows from Theorem 3.3. We recall a very well known result that a set of $n$ monomials in $n$ variables are algebrically independent if and only if the incidence matrix has determinant different by zero. Since the incidence matrix of a graph of monomials is the gradient matrix evaluated in $u_1 = 1, \dots, u_m = 1$, the first and the second case easily follow. In the last case $n > m$ hence the $g_i$ are algebrically dependent. The result follows from the Hessian criterion, Theorem 2.10 and the Gordan-Noether, Proposition 2.11. □

**Corollary 4.12.** *For all $r \geq 8$ there exist standard graded Artinian Gorenstein $\mathbb{K}$-algebras of socle degree 3 and $\operatorname{codim} A = r$, presented by quadrics, such that they not satisfy the WLP.*

*Proof.* For the second one, if $r = 9, 11$ the result follows from Examples 4.8 and 4.9.
For all $r = 2q \geq 8$ start with the square and then attach leaves as in Lemma 4.5 and the result follows. For all $r = 2q + 1 \geq 13$ start with the hexagon together with the central diagonal and then attach leaves as in Lemma 4.5 and the result follows. □

**Lemma 4.13.** *Let $f \in \mathbb{K}[x_1, \dots, x_r]$ be a homogeneous polynomial of degree $\deg(f) = 2k-1$ and such that $\operatorname{hess}_f^{k-1} = 0$ and let $\tilde{f} = uvf \in \mathbb{K}[x_1, \dots, x_n, u, v]$. Then $\operatorname{hess}_{\tilde{f}}^{k} = 0$.*

*Proof.* Let $R = \mathbb{K}[x_1, \dots, x_r]$ and $\tilde{R} = \mathbb{K}[x_1, \dots, x_r, u, v]$ and let $Q$ and $\tilde{Q}$ be the associated rings of differential operators. Let $A = Q/\operatorname{Ann}_Q(f)$ and let $\tilde{A} = \tilde{Q}/\operatorname{Ann}_{\tilde{Q}}(f)$. Applying twice Lemma 4.2, we get

$$\tilde{A}_k = A_{k-2}UV \oplus A_{k-1}U \oplus A_{k-1}V \oplus A_k.$$

Therefore,

$$\operatorname{Hess}_{\tilde{f}}^{k} = \begin{bmatrix} 0 & 0 & 0 & * \\ 0 & 0 & \operatorname{Hess}_f^k & * \\ 0 & \operatorname{Hess}_f^{k-1} & 0 & * \\ * & * & * & 0 \end{bmatrix}.$$

Using the Laplace expansion it is easy to check that $\operatorname{hess}_f^{k-1} = 0 \Rightarrow \operatorname{hess}_{\tilde{f}}^{k} = 0$.

□

**Corollary 4.14.** *There exists standard graded Artin Gorenstein algebras presented by quadrics of socle degree $d = 2k+1 \geq 3$ that do not satisfy the weak Lefschetz property when $\operatorname{codim} A \geq d + 5$. Furthermore, $A$ can be choosen to have Hilbert vector $\operatorname{Hilb}(A) = \sigma^{(d-3)}(1, r, r, 1)$.*

*Proof.* By Corollary 4.12, for all $r \geq 8$ exist $f_r \in R = \mathbb{K}[x_1, \dots, x_r]_3$ such that $\operatorname{hess}_{f_r} = 0$ and $A_r$ is presented by quadrics. Let

$$\tilde{f} = \tilde{f}_{r,k} = f_r u_1 \dots u_{2k-2} \in \tilde{R} = \mathbb{K}[x_1, \dots, x_r, u_1, \dots, u_{2k-2}].$$

We have $\deg \tilde{f} = 2k + 1 \geq 3$. Let $\tilde{A} = \tilde{Q}/\operatorname{Ann}_{\tilde{Q}}(\tilde{f})$. Then

$$\operatorname{codim} \tilde{A} = 2k - 2 + r \geq 2k + 6 = d + 5.$$

By Lemma 4.1, since $A_r$ is presented by quadrics, $\tilde{A}$ is also presented by quadrics.

By Lemma 4.2 and by induction on $k$, we get that the Hilbert vector of $\tilde{A}$ are maximal.

By induction on $k$ and Lemma 4.13, $\operatorname{hess}^k(\tilde{f}) = 0$, hence, by the Strong Lefschetz Hessian criterion,Theorem 2.10, $\tilde{A}$ does not have the WLP. □

4.3. **Artinian Gorenstein algebras with even socle degree.** Notice that Lemma 4.2 and Lemma 4.1 together with the Hessian criterion (Theorem 2.10) and the inductive procedure of Corollary 4.4 allow us to produce for any socle degree $d \geq 3$ and $\operatorname{codim} A \geq d+5$ Artinian Gorenstein algebras presented by quadrics that do not satisfy the SLP, but this construction, in even socle degree is not enough to fail the WLP.

**Remark 4.15.** For $f \in \mathbb{K}[x_1, \ldots, x_n, u_1, \ldots, u_m]_{(1,3)}$ be a quartic bihomogeneous polynomial of bidegree $(1,3)$ of monomial square free type. By Lemma 4.10 and by Lemma 3.6, $\operatorname{rk} \operatorname{Hess}_f^{(1,2)}$ is maximal if and only if $\operatorname{rk} \operatorname{Hess}_f^{((1,0),(0,2))} = n$.

**Example 4.16.** Let $A$ be the algebra associated to the complex $\mathcal{TK}(2,2,3) \setminus e$ where $e$ is an edge having two incident faces, it has codimension 17 and it does not have the WLP. Indeed, $|V| = 7$, $|E| = 15$ and $|F| = 10$ but the matrix $\operatorname{Hess}_f^{((0,1),(2,0))}$ does not have maximal rank.

**Corollary 4.17.** *For all $r \geq 16$ there exist standard graded Artinian Gorenstein $\mathbb{K}$-algebras of socle degree 4 and $\operatorname{codim} A = r$, presented by quadrics and not satisfying the WLP.*

*Proof.* For codimension $r = 2q \geq 14$, start with the Turan algebra $TA(2,2,2)$ of codimension 14, which by Corollary 3.9 does not have the WLP and attach leaves as in Lemma 4.5 and the result follows. For codimension $r = 2q+1 \geq 17$, start with the Turan algebra $TA(2,2,3)$ of codimension 17, which by Corollary 3.9 does not have the WLP and attach leaves to conlude the result. □

**Lemma 4.18.** *Let $A = Q/\operatorname{Ann}_Q(f)$ be a standard graded Artinian Gorenstein algebra of socle degree $2q$ with $f \in \mathbb{K}[x_1, \ldots, x_n]$ and let $\tilde{f} \in \mathbb{K}[x_1, \ldots, x_n, u, v]$ be $\tilde{f} = fuv$. If $\operatorname{rk} \operatorname{Hess}_f^{(q-1,q)} < \dim A_{q-1}$, then $\operatorname{rk} \operatorname{Hess}_{\tilde{f}}^{(q,q+1)} < \dim A_q$.*

*Proof.* Let $R = \mathbb{K}[x_1, \ldots, x_r]$ and $\tilde{R} = \mathbb{K}[x_1, \ldots, x_r, u, v]$ and let $Q, \tilde{Q}, A\tilde{A}$ as usual. Applying twice Lemma 4.2, we get

$$\tilde{A}_k = A_{k-2}UV \oplus A_{k-1}U \oplus A_{k-1}V \oplus A_k.$$

Therefore,

$$\operatorname{Hess}_{\tilde{f}}^{(q,q+1)} = \begin{bmatrix} 0 & 0 & 0 & \operatorname{Hess}_f^{(q-2,q+1)} \\ 0 & 0 & \operatorname{Hess}_f^{(q-1,q)} & * \\ 0 & \operatorname{Hess}_f^{(q-1,q)} & 0 & * \\ \operatorname{Hess}_f^{(q,q-1)} & * & * & 0 \end{bmatrix}.$$

The result follows by the shape of the matrix by simple inspection. The details are left to the reader.

□

**Corollary 4.19.** *There exists a standard graded Artinian Gorenstein algebras presented by quadrics of socle degree $d = 2k \geq 4$ that does not have the weak Lefschetz property for each codimension $\operatorname{codim} A \geq d + 12$.*

*Proof.* The result follows by the inductive procedure in the same way as Corollary 4.14. □

**Acknowledgments**. We wish to thank Francesco Russo for his suggestions and conversations on the subject and the participants to the Commutative Algebra/Algebraic Geometry Seminar of the *Università di Catania*. The first author also thanks the Sicilian hospitality he found everywhere in Catania, rendering his long staying there very pleasant and fruitful. The first author was Partially supported by the CAPES postdoctoral fellowship, Proc. BEX 2036/14-2. The second author is part of the Research Project of the University of Catania FIR 2014 "Aspetti geometrici e algebrici della Weak e Strong Lefschetz Property".

## References

[Co] A.Conca, *Grobner bases for spaces of quadrics of low codimension*, Adv. in Appl. Math. 24 (2000), no. 2, 111–124.

[CRS] C. Ciliberto, F. Russo, A. Simis, *Homaloidal hypersurfaces and hypersurfaces with vanishing Hessian*, Adv. in Math. 218 (2008), 1759-1805.

[DP] A. Dimca, S. Papadima, *Hypersurfaces complements, Milnor fibres and higher homotopy groups of arrangements*, Annals of Math. 158 (2003), 473-507.

[Go] R. Gondim *On higher Hessians and the Lefschetz properties*, arXiv:1506.06387.

[GN] P. Gordan, M. Nöther, *Ueber die algebraischen Formen, deren Hesse'sche Determinante identisch verschwindet*, Math. Ann. 10 (1876), 547–568.

[GR] A. Garbagnati, F. Repetto, *A geometrical approach to Gordan–Noether's and Franchetta's contributions to a question posed by Hesse*, Collect. Math. 60 (2009), 27–41.

[GRu] R. Gondim, F. Russo, *Cubic hypersurfaces with vanishing Hessian*, Journal of Pure and Applied Algebra 219 (2015), 779-806.

[He] O. Hesse, *Zur Theorie der ganzen homogenen Functionen*, J. reine angew. Math. 56 (1859), 263–269.

[HMNW] T. Harima, J. Migliore, U. Nagel and J. Watanabe, *The weak and strong Lefschetz properties for artinian K-algebras*, J. Algebra 262 (2003), 99 126.

[HMMNWW] T. Harima, T. Maeno, H. Morita, Y. Numata, A. Wachi, J. Watanabe, *The Lefschetz properties*, Lecture Notes in Mathematics 2080. Springer, Heidelberg, 2013, xx+250 pp.

[La] K.Lamotke *Topology of complex algebraic Varieties after S. Lefschetz*, Topology 20 (1981), 15–51.

[Lo] C. Lossen, *When does the Hessian determinant vanish identically? (On Gordan and Noether's Proof of Hesse's Claim)*, Bull. Braz. Math. Soc. 35 (2004), 71–82.

[MN1] J. Migliore, U. Nagel *Survey article: a tour of the weak and strong Lefschetz properties*, J. Commut. Algebra 5 (2013), no. 3, 329–358.

[MN2] J. Migliore, U. Nagel *Gorenstein algebras presented by quadrics*, Collect. Math. 62 (2013), 211–233.

[MW] T. Maeno, J. Watanabe, *Lefschetz elements of artinian Gorenstein algebras and Hessians of homogeneous polynomials*, Illinois J. Math. 53 (2009), 593–603.

[Pe] U. Perazzo, *Sulle varietà cubiche la cui hessiana svanisce identicamente*, Giornale di Matematiche (Battaglini) 38 (1900), 337–354.

[Ru] F. Russo *On the Geometry of Some Special Projective Varieties*, Lecture Notes of the Unione Matematica Italiana, vol. 18 Springer (2016).

[St] R.Stanley, *Weyl groups, the hard Lefschetz theorem, and the Sperner property*, SIAM J. Algebraic Discrete Methods 1 (1980), 168–184.

[St2] R. Stanley, *Hilbert functions of graded algebras*, Adv. in Math. 28 (1978), 57–83

[Tu] Turán, Paul (1941), *On an extremal problem in graph theory*, Matematikai és Fizikai Lapok (in Hungarian) 48: 436–452.

[Wa1] J. Watanabe, *A remark on the Hessian of homogeneous polynomials*, in *The Curves Seminar at Queen's*, Volume XIII, Queen's Papers in Pure and Appl. Math. 119 (2000), 171–178.

[Wa2] J. Watanabe, *On the Theory of Gordan-Noether on Homogeneous Forms with Zero Hessian* , Proc. Sch. Sci. TOKAI UNIV. 49 (2014), 1–21.

Universidade Federal Rural de Pernambuco, av. Don Manoel de Medeiros s/n, Dois Irmãos - Recife - PE 52171-900, Brasil
*E-mail address*: rodrigo.gondim@ufrpe.br

Dipartimento di Matematica e Informatica, Università degli Studi di Catania, Viale A. Doria 5, 95125 Catania, Italy
*E-mail address*: zappalag@dmi.unict.it